

## INDEX TO VOLUME 8

### AUTHORS

- AESHITA, M., B. WILSKE, T. ZHAN-HUI, and J. CHEN  
Occurrence and morphometric variability in the frugivorous bat species, *Cynopterus sphinx* and *Rousettus leschenaulti*, from a tropical rain forest, Xishuangbanna, SW-China, 417
- AIHARTZA, J. R.  
see ALMENAR, D., 157  
see GOITI, U., 141
- ALMENAR, D., J. AIHARTZA, U. GOITI, E. SALSAMENDI, and I. GARIN  
Habitat selection and spatial use by the trawling bat *Myotis capaccinii* (Bonaparte, 1837), 157
- ALMENAR, D.  
see GOITI, U., 141
- ANDREWS, M. M., P. T. ANDREWS, D. F. WILLS, and S. M. BEVIS  
Ultrasound social calls of greater horseshoe bats (*Rhinolophus ferrumequinum*) in a hibernaculum, 197
- ANDREWS, P. T.  
see ANDREWS, M. M., 197
- ANDRIAFIDISON, D.  
see RANDRIANANDRIANINA, F., 429
- ANDRIANAIVOARIVELO, A. R., N. RANAIVOSON, P. A. RACEY, and R. K. B. JENKINS  
The diet of three synanthropic bats (Chiroptera: Molossidae) from eastern Madagascar, 439
- ARMSTRONG, K. N.  
Phylogeographic structure in *Rhinonictis aurantia* (Chiroptera: Hipposideridae): implications for conservation, 63
- BARCLAY, L. E.  
see BARCLAY, R. M. R., 549
- BARCLAY, R. M. R., L. E. BARCLAY, and D. S. JACOBS  
Deliberate insectivory by the fruit bat *Rousettus aegyptiacus*, 549
- BARNETT, A. A., E. M. SAMPAIO, E. K. V. KALKO, R. L. SHAPLEY, E. FISCHER, G. CAMARGO, and B. RODRÍGUEZ-HERRERA  
Bats of Jaú National Park, central Amazônia, Brazil, 103
- BASHTA, A.-T.  
see SACHANOWICZ, K., 543
- BATES, P. J. J., F. H. RATRIMOMANARIVO, D. L. HARRISON, and S. M. GOODMAN  
A description of a new species of *Pipistrellus* (Chiroptera: Vespertilionidae) from Madagascar with a review of related Vespertilioninae from the island, 299
- BATES, P. J. J.  
see BUMRUNGSRI, S., 325  
see THONG, V. D., 83
- BENDA, P.  
see JONES, G., 451
- BERKOVÁ, H., and J. ZUKAL  
Flight activity of bats at the entrance of a natural cave, 187
- BEVIS, S. M.  
see ANDREWS, M. M., 197
- BRINKMANN, R.  
see STECK, C. E., 445
- BUMRUNGSRI, S., D. L. HARRISON, C. SATASOOK, A. PRAJUKJITR, S. THONG-AREE, and P. J. J. BATES  
A review of bat research in Thailand with eight new species records for the country, 325
- BUMRUNGSRI, S.  
see THONG V. D., 83
- CAMARGO, G.  
see BARNETT, A. A., 103
- CARTER, G. G., C. E. COEN, L. M. STENZLER, and I. J. LOVETTE  
Avian host DNA isolated from the feces of white-winged vampire bats (*Diaemus youngi*), 255
- CHATTOPADHYAY, B.  
see RAJASEKAR, R., 269
- CHEN, J.  
see AESHITA, M., 417
- COEN, C. E.  
see CARTER, G. G., 255
- CSORBA, G.  
see KUO, H.-C., 11
- DENZINGER, A.  
see DIETRICH, S., 465
- DEWA, H.  
see KAWAI, K., 95
- DIETRICH, S., D. P. SZAMEITAT, A. KIEFER, H.-U.

- SCHNITZLER, and A. DENZINGER  
Echolocation signals of the plecotine bat, *Plecotus macrobullaris* Kuzyakin, 1965, 465
- DIETZ, C., I. DIETZ, T. IVANOVA, and B. M. SIEMERS  
Effects of forearm bands on horseshoe bats (Chiroptera: Rhinolophidae), 523
- DIETZ, I.  
see DIETZ, C., 523
- DIETZ, M., J. A. ENCARNAÇÃO, and E. K. V. KALKO  
Small scale distribution patterns of female and male Daubenton's bats (*Myotis daubentonii*), 403
- DOWNS, N. C., and P. A. RACEY  
The use by bats of habitat features in mixed farmland in Scotland, 169
- DOWNS, N. C.  
see PETRŽELKOVÁ, K. J., 381
- ELANGOVA, V., E. Y. S. PRIYA, and G. MARIMUTHU  
Olfactory discrimination ability of the short-nosed fruit bat *Cynopterus sphinx*, 247
- ENCARNAÇÃO, J. A.  
see DIETZ, M., 403
- FANG, Y.-P.  
see KUO, H.-C., 11
- FENTON, M. B.  
see SCHMIDT-FRENCH, B., 477
- FISCHER, E.  
see BARNETT, A. A., 103
- FRANCIS, C. M.  
see GUILLÉN-SERVENT, A., 39
- FUKUI, D.  
see KAWAI, K., 95
- GAMBA-RÍOS, M.  
see RODRÍGUEZ-HERRERA, B., 557
- GARCÍA-MUDARRA, J. L.  
see IBÁÑEZ, C., 287
- GARIN, I.  
see GOITI, U., 141  
see ALMENAR, D., 157
- GERLACH, J., and M. TAYLOR  
Habitat use, roost characteristics and diet of the Seychelles sheath-tailed bat *Coleura seychelensis*, 129
- GILLAM, E.  
see SCHMIDT-FRENCH, B., 477
- GOITI, U., J. R. AIHARTZA, D. ALMENAR, E. SALSAMENDI, and I. GARIN  
Seasonal foraging by *Rhinolophus euryale* (Rhinolophidae) in an Atlantic rural landscape in northern Iberian Peninsula, 141
- GOITI, U.  
see ALMENAR, D., 157
- GOODMAN, S. M., and O. GRIFFITHS  
A case of exceptionally high predation levels of *Rousettus madagascariensis* by *Tyto alba* (Aves: Tytonidae) in western Madagascar, 553
- GOODMAN, S. M., F. H. RATRIMOMANARIVO, and F. H. RANDRIANANDRIANINA  
A new species of *Scotophilus* (Chiroptera: Vespertilionidae) from western Madagascar, 21
- GOODMAN, S. M.  
see BATES, P. J. J., 299
- GRIFFITHS, O.  
see GOODMAN, S. M., 553
- GUILLÉN-SERVENT, A., and C. M. FRANCIS  
A new species of bat of the *Hipposideros bicolor* group (Chiroptera: Hipposideridae) from Central Laos, with evidence of convergent evolution with Sundaic taxa, 39
- HARBUSCH, C., and P. A. RACEY  
The sessile serotine: the influence of roost temperature on philopatry and reproductive phenology of *Eptesicus serotinus* (Schreber, 1774) (Mammalia: Chiroptera), 213
- HARRISON, D. L.  
see BATES, P. J. J., 299  
see BUMRUNGSI, S., 325  
see THONG V. D., 83
- HELGEN, K. M.  
see THONG V. D., 83
- HELVENSEN, O. VON  
see SACHTELEBEN, J., 391
- HIRAKAWA, H., and K. KAWAI  
Hiding low in the thicket: roost use by Ussurian tube-nosed bats (*Murina ussuriensis*), 263
- HOCKING, P.  
see PACHECO, V., 561
- HORÁČEK, I.  
see HULVA, P., 537
- HULVA, P., and I. HORÁČEK  
Cryptic sympatric diversity in *Emballonura alecto*: further bat species?, 537
- HUTCHEON, J. M., and J. A. W. KIRSCH  
A moveable face: deconstructing the Microchiroptera and a new classification of extant bats, 1
- IBÁÑEZ, C., J. L. GARCÍA-MUDARRA, M. RUEDI, B. STADELMANN, and J. JUSTE  
The Iberian contribution to cryptic diversity in European bats, 287
- IVANOVA, T.  
see DIETZ, C., 523
- JACOBS, D. S.  
see BARCLAY, R. M. R., 549
- JENKINS, R. K. B.  
see ANDRIANAIVOARIVELO, A. R., 439  
see RANDRIANANDRIANINA, F., 429
- JONES, G., S. PARSONS, S. ZHANG, B. STADELMANN, P. BENDA, and M. RUEDI  
Echolocation calls, wing shape, diet and phylogenetic diagnosis of the endemic Chinese bat *Myotis pequinus*, 451

- JUSTE, J.  
see IBÁÑEZ, C., 287
- KALKO, E. K. V.  
see BARNETT, A. A., 103  
see DIETZ, M., 403
- KAWAI, K., N. KONDO, N. SASAKI, D. FUKUI, H. DEWA, M. SATÔ, and Y. YAMAGA  
Distinguishing between cryptic species *Myotis ikonnikovi* and *M. brandtii gracilis* in Hokkaido, Japan: evaluation of a novel diagnostic morphological feature using molecular methods, 95
- KAWAI, K.  
see HIRAKAWA, H., 263
- KIEFER, A.  
see DIETRICH, S., 465
- KIRSCH, J. A. W.  
see HUTCHEON, J. M., 1
- KOFOKY, A. F.  
see RANDRIANANDRIANINA, F., 429
- KOKUREWICZ, T., and J. R. SPEAKMAN  
Age related variation in the energy costs of torpor in Daubenton's bat: effects on fat accumulation prior to hibernation, 509
- KONDO, N.  
see KAWAI, K., 95
- KRUTZSCH, P. H., and D. W. NELLIS  
Reproductive anatomy and cyclicity of the male bat *Brachyphylla cavernarum* (Chiroptera: Phyllostomidae), 497
- KUO, H.-C., Y.-P. FANG, G. CSORBA, and L.-L. LEE  
The definition of *Harpiola* (Vespertilionidae: Muriniinae) and the description of a new species from Taiwan, 11
- LEE, L.-L.  
see KUO, H.-C., 11
- LOVETTE, I. J.  
see CARTER, G. G., 255
- MALDONADO, J. E.  
see ORTEGA, J., 485
- MARIMUTHU, G.  
see ELANGOVAN, V., 247  
see SINGARAVELAN, N., 239
- MEDELLÍN, R. A.  
see RODRÍGUEZ-HERRERA, B., 557
- MUÑOZ-ROMO, M.  
Ethogram and diurnal activities of a colony of *Artibeus lituratus* (Phyllostomidae: Stenodermatinae), 231
- NELLIS, D. W.  
see KRUTZSCH, P. H., 497
- ORTEGA, J., and J. E. MALDONADO  
Female interactions in harem groups of the Jamaican fruit-eating bat, *Artibeus jamaicensis* (Chiroptera: Phyllostomidae), 485
- PACHECO, V., and P. HOCKING  
Notably range extension of *Sturnira aratathomasi* Peterson and Tamsitt 1969 in Perú, 561
- PARSONS, S.  
see JONES, G., 451
- PEARCE, M. J.  
see THONG V. D., 83
- PETRŽELKOVÁ, K. J., N. C. DOWNS, J. ZUKAL, and P. A. RACEY  
A comparison between emergence and return activity in pipistrelle bats *Pipistrellus pipistrellus* and *P. pygmaeus*, 381
- PIKSA, K.  
Aggressive behaviour of greater mouse-eared bat (*Myotis myotis*) towards lesser horseshoe bats (*Rhinolophus hipposideros*) in a hibernaculum, 566
- PRAJUKIJTR, A.  
see BUMRUNGSI, S., 325
- PRIYA, E. Y. S.  
see ELANGOVAN, V., 247
- RACEY, P. A.  
see ANDRIANAIVOARIVELO, A. R., 439  
see DOWNS, N. C., 169  
see HARBUSCH, C., 213  
see PETRŽELKOVÁ, K. J., 381  
see RANDRIANANDRIANINA, F., 429
- RAJASEKAR, R., B. CHATTOPADHYAY, and K. SRIPATHI  
Depositing masticated plant materials inside tent roosts in *Cynopterus sphinx* (Chiroptera: Pteropodidae) in Southern India, 269
- RAMILJAONA, O.  
see RANDRIANANDRIANINA, F., 429
- RANAIVOSON, N.  
see ANDRIANAIVOARIVELO, A. R., 439
- RANDRIANANDRIANINA, F., D. ANDRIAFIDISON, A. F. KOFOKY, O. RAMILJAONA, F. RATRIMOMANARIVO, P. A. RACEY, and R. K. B. JENKINS  
Habitat use and conservation of bats in rainforest and adjacent human-modified habitats in eastern Madagascar, 429
- RANDRIANANDRIANINA, F. H.  
see GOODMAN, S. M., 21
- RATRIMOMANARIVO, F.  
see RANDRIANANDRIANINA, F., 429
- RATRIMOMANARIVO, F. H.  
see BATES, P. J. J., 299  
see GOODMAN, S. M., 21
- RITZI, C. M.  
see WALTERS, B. L., 259
- RODRÍGUEZ-HERRERA, B., R. A. MEDELLÍN, and M. GAMBA-RÍOS  
Tent building by female *Ectophylla alba* (Chiroptera: Phyllostomidae) in Costa Rica, 557
- RODRÍGUEZ-HERRERA, B.  
see BARNETT, A. A., 103

- RUEDI, M.  
 see IBÁÑEZ, C., 287  
 see JONES, G., 451
- SACHANOWICZ, K., A. WOWER, and A.-T. BASHTA  
 Further range extension of *Pipistrellus kuhlii* (Kuhl, 1817) in central and eastern Europe, 543
- SACHTELEBEN, J., and O. VON HELVERSEN  
 Songflight behaviour and mating system of the pipistrelle bat (*Pipistrellus pipistrellus*) in an urban habitat, 391
- SALSAMENDI, E.  
 see ALMENAR, D., 157  
 see GOITI, U., 141
- SAMPAIO, E. M.  
 see BARNETT, A. A., 103
- SASAKI, N.  
 see KAWAI, K., 95
- SATASOOK, C.  
 see BUMRUNGSRI, S., 325
- SATÔ, M.  
 see KAWAI, K., 95
- SCHMIDT-FRENCH, B., E. GILLAM, and M. B. FENTON  
 Vocalizations emitted during mother-young interactions by captive eastern red bats *Lasiurus borealis* (Chiroptera: Vespertilionidae), 477
- SCHNITZLER, H.-U.  
 see DIETRICH, S., 465
- SHAPLEY, R. L.  
 see BARNETT, A. A., 103
- SIEMERS, B. M.  
 see DIETZ, C., 523
- SINGARAVELAN, N., and G. MARIMUTHU  
*Muntingia calabura* — an attractive food plant of *Cynopterus sphinx* — deserves planting to lessen orchard damage, 239
- SPARKS, D. W.  
 see WALTERS, B. L., 259
- SPEAKMAN, J. R.  
 see KOKUREWICZ, T., 509
- SRIPATHI, K.  
 see RAJASEKAR, R., 269
- STADELMANN, B.  
 see IBÁÑEZ, C., 287  
 see JONES, G., 451
- STECK, C. E., and R. BRINKMANN  
 The trophic niche of the Geoffroy's bat (*Myotis emarginatus*) in south-western Germany, 445
- STENZLER, L. M.  
 see CARTER, G. G., 255
- SZAMEITAT, D. P.  
 see DIETRICH, S., 465
- TAYLOR, M.  
 see GERLACH, J., 129
- TEJEDOR, A.  
 The type locality of *Natalus stramineus* (Chiroptera: Natalidae): implications for the taxonomy and biogeography of the genus *Natalus*, 361
- THONG V. D., S. BUMRUNGSRI, D. L. HARRISON, M. J. PEARCH, K. M. HELGEN, and P. J. J. BATES  
 New records of Microchiroptera (Rhinolophidae and Kerivoulinae) from Vietnam and Thailand, 83
- THONG-AREE, S.  
 see BUMRUNGSRI, S., 325
- WALTERS, B. L., D. W. SPARKS, J. O. WHITAKER, JR., and C. M. RITZI  
 Timing of migration by eastern red bats (*Lasiurus borealis*) through Central Indiana, 259
- WHITAKER, J. O., JR.,  
 see WALTERS, B. L., 259
- WILLS, D. F.  
 see ANDREWS, M. M., 197
- WILSKE, B.  
 see AESHITA, M., 417
- WOWER, A.  
 see SACHANOWICZ, K., 543
- YAMAGA, Y.  
 see KAWAI, K., 95
- ZHANG, S.  
 see JONES, G., 451
- ZHAN-HUI, T.  
 see AESHITA, M., 417
- ZUKAL, J.  
 see BERKOVÁ, H., 187  
 see PETRŽELKOVÁ, K. J., 381